

WEST Search History

DATE: Thursday, February 23, 2006

Hide?	<u>Set</u> <u>Name</u>	<u>Query</u>	<u>Hit</u> <u>Count</u>
	<i>DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI,TDBD; PLUR=YES; OP=OR</i>		
<input type="checkbox"/>	L1	blood near group near binding	15
<input type="checkbox"/>	L2	hops or hop-s or hops	41251
<input type="checkbox"/>	L3	L2 and (helicobacter or pylori)	70
<input type="checkbox"/>	L4	L2 same (helicobacter or pylori)	15
<input type="checkbox"/>	L5	outer near membrane near (polypeptide or peptide or protein)	3608
<input type="checkbox"/>	L6	L5 and (pylori or pyloridis or pylon or pylorum or pyloris or helicobacter or heliobacter)	534
<input type="checkbox"/>	L7	L5.clm. and (pylori or pyloridis or pylon or pylorum or pyloris or helicobacter or heliobacter).clm.	11
<input type="checkbox"/>	L8	L7 not l1	11
<input type="checkbox"/>	L9	L8 not l4	10
<input type="checkbox"/>	L10	bab or baba or bab-a or baba1 or baba2 or bab-a or babb or bab-b or babc or bab-c or bab-a1 or bab-a2	35558
<input type="checkbox"/>	L11	L10 and (helicobacter or pylori or pyroli or pyloridis or pylorid or pyloridis or hpylori)	143
<input type="checkbox"/>	L12	L10.clm. and (helicobacter or pylori or pyroli or pyloridis or pylorid or pyloridis or hpylori)	5
<input type="checkbox"/>	L13	73.clm. or 74.clm. or 75.clm. or 76.clm.	100680
<input type="checkbox"/>	L14	L13 same (heliobacter or helicobacter or pylori)	31

END OF SEARCH HISTORY

DOCUMENT-IDENTIFIER: US 20050009037 A1

TITLE: *Helicobacter bizzozeronii* outer membrane protein encoding gene and its use in diagnostic and treatment methods

Summary of Invention Paragraph:

[0007] Several families of outer-membrane protein (Omp) have been identified in *H. pylori* (Alm et al., "Comparative Genomics of *Helicobacter pylori*: Analysis of the Outer Membrane Protein Families," Infect. Immun. 68:4155-4168 (2000)). One major family (*Hop*) contains thirty-two homologous members, some of which are functional as porins (Peck et al., "Characterization of Four Members of a Multigene Family Encoding Outer Membrane Proteins of *Helicobacter pylori* and Their Potential for Vaccination," Microbes Infect. 3:171-179 (2001)). Some outer-membrane proteins not in paralogous families have been also identified. Among them are small-molecular-mass bacterial peptidoglycan-associated lipoproteins (PAL) such as Lpp20 and Omp22 (Kostrzynska et al., "Molecular Characterization of a Conserved 20-Kilodalton Membrane-Associated Lipoprotein Antigen of *Helicobacter pylori*," J. Bacteriol. 176:5938-5948 (1994); Kim et al., "Cloning and Characterization of a 22 kDa Outer-Membrane Protein (Omp22) from *Helicobacter pylori*," Mol. Cells 10:633-641 (2000)).

WEST Search History

[Hide Items](#)[Restore](#)[Clear](#)[Cancel](#)

DATE: Thursday, February 23, 2006

Hide?	<u>Set Name</u>	<u>Query</u>	<u>Hit Count</u>
		<i>DB=USPT; PLUR=YES; OP=OR</i>	
<input type="checkbox"/>	L1	human.clm. same colostrum.clm.	18
<input type="checkbox"/>	L2	sig.a.clm. or s-iga.clm.	6
<input type="checkbox"/>	L3	L2 and colostrum	0
<input type="checkbox"/>	L4	secretory near3 iga	426
<input type="checkbox"/>	L5	L4 and colostrum	68
<input type="checkbox"/>	L6	l5 and (man or human or sapian or sapien)	67
<input type="checkbox"/>	L7	l5 and ((man or human or sapian or sapien) same colostrum)	38

END OF SEARCH HISTORY

CLUSTAL W (1.81) multiple sequence alignment

```

unk|VIRT5690|Blast_submission  -----EDDGFYTSVGYQIGEEAAQMV-----
tr|Q9ZMK5|Q9ZMK5_HELPJ      MKKTILLSLMVSSLFAENDGVYMSVGYQIGEEAAQMVKNTGEIQKVSNA
                               *:*. * *****

unk|VIRT5690|Blast_submission  -----
tr|Q9ZMK5|Q9ZMK5_HELPJ      NLNNLLTRYNELKQTASNTDSSTAQAIDNLEKSASRLKTTPTNTANQAVS

unk|VIRT5690|Blast_submission  -----
tr|Q9ZMK5|Q9ZMK5_HELPJ      ALSSAVGMWQVIASNLANNLSLSSEYEKCLKATSQLLQNTLENKNNNLKI

unk|VIRT5690|Blast_submission  -----
tr|Q9ZMK5|Q9ZMK5_HELPJ      NDYDQLLTQASTIINTLQSQCPGVDGGNGKPWGINTSGNACAIFGSTFN

unk|VIRT5690|Blast_submission  -----
tr|Q9ZMK5|Q9ZMK5_HELPJ      INSMIDSAKKAADARRTAPESPNQQNAFTNADFNKNLNQVSSVINDTI

unk|VIRT5690|Blast_submission  -----
tr|Q9ZMK5|Q9ZMK5_HELPJ      YLKGDNLETIYNTIQKTPNSKGFQSLVSRSSYSYSLNETQYSQFQTTTK

unk|VIRT5690|Blast_submission  -----
tr|Q9ZMK5|Q9ZMK5_HELPJ      FGHNPFPSVGLINSQSNNGAMNGVGVQLGYKQFFGKNKFFGIRYYGFFD

unk|VIRT5690|Blast_submission  -----
tr|Q9ZMK5|Q9ZMK5_HELPJ      NYAYIKSNFFNSASNVFTYAGSDLLLNFINGGSDRNRKVSFGIFGGIA

unk|VIRT5690|Blast_submission  -----
tr|Q9ZMK5|Q9ZMK5_HELPJ      AGTTWLNNQSANLKITNSAYSAKINNTNFQFLFNTGLRLQGIHHGIELG

unk|VIRT5690|Blast_submission  -----
tr|Q9ZMK5|Q9ZMK5_HELPJ      KIPTINTNYYSFMGAKLAYRRLYSLYLNVLAY



```

<input type="checkbox"/>	tr Q9ZKV2	_HELPE	Outer membrane protein-adhesin [babA]	[Helicobac
<input type="checkbox"/>	tr O25840	_HELPE	Outer membrane protein (Omp28) [HP1243]	[Helicob
<input type="checkbox"/>	tr O25556	_HELPE	Outer membrane protein (Omp19) [HP0896]	[Helicob
<input type="checkbox"/>	tr O25086	_HELPE	Outer membrane protein (Omp9) [HP0317]	[Helicoba
<input type="checkbox"/>	tr Q8GNW2	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNW1	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNW0	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNV8	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNV7	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNV6	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q7WVA4	_HELPE	Adhesin-binding fucosylated histo-blood group an	
<input type="checkbox"/>	tr Q7WVA2	_HELPE	Adhesin-binding fucosylated histo-blood group an	
<input type="checkbox"/>	tr Q7WVA1	_HELPE	Adhesin-binding fucosylated histo-blood group an	
<input type="checkbox"/>	tr Q7WV99	_HELPE	Adhesin-binding fucosylated histo-blood group an	
<input type="checkbox"/>	tr Q7WV84	_HELPE	Adhesin-binding fucosylated histo-blood group an	
<input type="checkbox"/>	tr Q7WV81	_HELPE	Adhesin-binding fucosylated histo-blood group an	
<input type="checkbox"/>	tr Q7WV74	_HELPE	Adhesin-binding fucosylated histo-blood group an	
<input type="checkbox"/>	tr Q7WV69	_HELPE	Adhesin-binding fucosylated histo-blood group an	
<input type="checkbox"/>	tr Q6T8D5	_HELPE	BabB (BabB2) [Helicobacter pylori (Campylobacter	
<input type="checkbox"/>	tr Q6T8D3	_HELPE	BabA [Helicobacter pylori (Campylobacter pylori)	
<input type="checkbox"/>	tr Q5Q1P2	_HELPE	BabB/BabA1 fusion protein 1 (Fragment) [babB/bab	
<input type="checkbox"/>	tr Q5Q1P1	_HELPE	BabB/BabA1 fusion protein 2 (Fragment) [babB/bab	
<input type="checkbox"/>	tr O52269	_HELPE	Adhesin binding fucosylated histo-blood group an	
<input type="checkbox"/>	tr O51811	_HELPE	Adhesin-binding fucosylated histo-blood group an	
<input type="checkbox"/>	tr Q6U2C5	_HELPE	Adhesin-binding fucosylated histo-blood group (F	
<input type="checkbox"/>	tr Q5G5J5	_HELPE	Adhesin-binding fucosylated histo-blood group (F	
<input type="checkbox"/>	tr Q8GNV9	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNY0	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNX9	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNX7	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNX5	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNX4	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNX3	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNX2	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNX1	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNW9	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNW8	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNW7	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNW6	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNW5	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNW4	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam
<input type="checkbox"/>	tr Q8GNW3	_HELPE	BabB (Fragment) [babB]	[Helicobacter pylori (Cam

<input type="checkbox"/>	tr Q7WVA5	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WVA0	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV97	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV96	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV95	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV94	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV93	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV92	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV90	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV89	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV88	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV87	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV86	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV85	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV83	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV82	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV80	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV79	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV75	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV70	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV68	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV67	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q7WV66	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q6JAA1	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q6JAA0	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q6JA99	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q6JA98	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q6JA97	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q5G5J6	_HELPY	Adhesin-binding	fucosylated	histo-blood	group (F
<input type="checkbox"/>	tr Q8GNX6	_HELPY	BabB (Fragment)	[babB]	[Helicobacter pylori (Cam	
<input type="checkbox"/>	tr Q7WV98	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr O34523	_HELPY	Outer membrane protein (Omp29)	(Omp5)	[HP0227]	[
<input type="checkbox"/>	tr Q6U2D4	_HELPY	Adhesin-binding	fucosylated	histo-blood	group (F
<input type="checkbox"/>	tr Q6U2D5	_HELPY	Adhesin-binding	fucosylated	histo-blood	group (F
<input type="checkbox"/>	tr Q5G5J4	_HELPY	Adhesin-binding	fucosylated	histo-blood	group (F
<input type="checkbox"/>	tr Q8GNX8	_HELPY	BabB (Fragment)	[babB]	[Helicobacter pylori (Cam	
<input type="checkbox"/>	tr Q7WVA3	_HELPY	Adhesin-binding	fucosylated	histo-blood	group an
<input type="checkbox"/>	tr Q6U2D3	_HELPY	Adhesin-binding	fucosylated	histo-blood	group (F
<input type="checkbox"/>	tr Q9Z390	_HELPJ	Putative Outer membrane protein	[jhp_0212]	[Heli	
<input type="checkbox"/>	tr Q9ZN51	_HELPJ	Putative Outer membrane protein	[jhp_0007]	[Heli	
<input type="checkbox"/>	tr Q9X748	_HELPY	HopZ protein precursor	[hopZ]	[Helicobacter pylo	
<input type="checkbox"/>	tr Q9X747	_HELPY	HopZ protein precursor	[hopZ]	[Helicobacter pylo	

☐ tr Q9X746 _HELPY HopZ protein precursor [hopZ] [Helicobacter pylo
☐ tr Q9X745 _HELPY HopZ protein precursor [hopZ] [Helicobacter pylo
☐ tr Q9S3I7 _HELPY HopZ protein precursor [hopZ] [Helicobacter pylo
☐ tr Q7X2J7 _HELPY HopZ (Fragment) [hopZ] [Helicobacter pylori (Cam
☐ tr Q7WV78 _HELPY Adhesin-binding fucosylated histo-blood group an
☐ tr Q7WV76 _HELPY Adhesin-binding fucosylated histo-blood group an
☐ tr Q7WV71 _HELPY Adhesin-binding fucosylated histo-blood group an
☐ tr Q9ZMK5 _HELPJ Outer membrane protein/porin [hopA] [Helicobacte
☐ tr Q6U2C8 _HELPY Adhesin-binding fucosylated histo-blood group (F
☐ tr Q8GNX0 _HELPY BabB (Fragment) [babB] [Helicobacter pylori (Cam
☐ tr Q7WV91 _HELPY Adhesin-binding fucosylated histo-blood group an
☐ tr Q7WV77 _HELPY Adhesin-binding fucosylated histo-blood group an
☐ tr Q7WV73 _HELPY Adhesin-binding fucosylated histo-blood group an
☐ tr Q7WV72 _HELPY Adhesin-binding fucosylated histo-blood group an
☐ tr Q9ZN38 _HELPJ Putative Outer membrane protein [jhp_0021] [Heli
☐ tr O24870 _HELPY Outer membrane protein (Omp2) [HP0025] [Helicoba

Graphical overview of the alignments

 to resubmit your query after masking regions matching PROSITE
profiles or Pfam HMMs
( Help) (use ScanProsite for more details about PROSITE matches)

Profile hits



Pfam hits



Submission	Matches on query sequence	
1		1
Q9ZKV2_HELPJ		0
Q25840_HELPY		0
Q25556_HELPY		0
Q25886_HELPY		0
Q8GN42_HELPY		0
Q8GN41_HELPY		0
Q8GN40_HELPY		0
Q8GNV8_HELPY		0
Q8GNV7_HELPY		0
Q8GNV6_HELPY		0
Q7HVR4_HELPY		0
Q7HVR2_HELPY		0
Q7HVR1_HELPY		0
Q7HV99_HELPY		0
Q7HV84_HELPY		0
Q7HV81_HELPY		0
Q7HV74_HELPY		0
Q7HV69_HELPY		0
Q6T805_HELPY		0
Q6T803_HELPY		0
Q5Q1P2_HELPY		0
Q5Q1P1_HELPY		0
Q52269_HELPY		0
Q51811_HELPY		0
Q6U2C5_HELPY		0
Q5G5J5_HELPY		0
Q8GNV9_HELPY		0
Q8GNY8_HELPY		0
Q8GNX9_HELPY		0
Q8GNX7_HELPY		0
Q8GNX5_HELPY		0
Q8GNX4_HELPY		0
Q8GNX3_HELPY		0
Q8GNX2_HELPY		0
Q8GNX1_HELPY		0
Q8GN49_HELPY		0
Q8GN48_HELPY		0
Q8GN47_HELPY		0
Q8GN46_HELPY		0
Q8GN45_HELPY		0
Q8GN44_HELPY		0
Q8GN43_HELPY		0
Q7HVR5_HELPY		0
Q7HVR8_HELPY		0
Q7HV97_HELPY		0
Q7HV96_HELPY		0
Q7HV95_HELPY		0
Q7HV94_HELPY		0
Q7HV93_HELPY		0
Q7HV92_HELPY		0
Q7HV90_HELPY		0
Q7HV89_HELPY		0
Q7HV88_HELPY		0
Q7HV87_HELPY		0
Q7HV86_HELPY		0
Q7HV85_HELPY		0
Q7HV83_HELPY		0
Q7HV82_HELPY		0
Q7HV80_HELPY		0
Q7HV79_HELPY		0
Q7HV75_HELPY		0
Q7HV70_HELPY		0
Q7HV68_HELPY		0
Q7HV67_HELPY		0
Q7HV66_HELPY		0
Q6JAA1_HELPY		0
Q6JAA0_HELPY		0
Q6JA99_HELPY		0
Q6JA98_HELPY		0
Q6JA97_HELPY		0
Q5G5J6_HELPY		0
Q8GN06_HELPY		0
Q7HV98_HELPY		0
Q34529_HELPY		0
Q6U2D4_HELPY		0
Q6U2D5_HELPY		0
Q5G5J4_HELPY		0
Q8GN08_HELPY		0
Q7HVR3_HELPY		0
Q6U2D3_HELPY		0
Q9Z988_HELPJ		0
Q9ZM51_HELPJ		0
Q9X748_HELPY		0
Q9X747_HELPY		0
Q9X746_HELPY		0

□ 1: FEMS Immunol Med Microbiol. 1994 Jun;9(1):15-21.

[Related Articles, Links](#)

Detection of microbial surface antigens that bind Lewis(a) antigen.

Essery SD, Weir DM, James VS, Blackwell CC, Saadi AT, Busuttil A, Tzanakaki G.

Department of Medical Microbiology, University Medical School, University of Edinburgh, UK.

There is evidence that the Lewis(a) blood group antigen is one of the receptors for a number of potentially pathogenic microorganisms. To determine how widely distributed the microbial adhesins are that bind this antigen, anti-idiotypic antibodies produced against monoclonal anti-Lewis(a) were used in coagglutination assays to screen a variety of species. The following were agglutinated: 7/7 strains of *Staphylococcus aureus*; 10/19 (53%) strains of *Neisseria meningitidis*; 8/13 (62%) strains of *Haemophilus influenzae*; 1/3 strains of *Helicobacter pylori*; 1/2 strains of *Neisseria gonorrhoeae*; 1/2 strains of *Candida albicans*. The application of the anti-idiotypic antibodies to studies of host cell receptors, isolation of adhesins and development of new epidemiological typing reagents is discussed.

PMID: 7920460 [PubMed - indexed for MEDLINE]

A handwritten signature, 'Adonis', is enclosed within a hand-drawn oval. The signature is written in a cursive, flowing style.

Gastroenterology. 1997 Apr;112(4):1179-87.

Related Articles, Links

Gastroenterology

Isolation of a cell surface component of *Helicobacter pylori* that binds H type 2, Lewis(a), and Lewis(b) antigens.

Alkout AM, Blackwell CC, Weir DM, Poxton IR, Elton RA, Luman W, Palmer K.

Department of Medical Microbiology, University of Edinburgh, Scotland.

BACKGROUND & AIMS: Individuals of blood group O and nonsecretors of ABO blood group antigens are more susceptible to peptic ulcers. The aim of this study was to determine if blood group antigens associated with group O or secretor status are epithelial cell receptors for *Helicobacter pylori*. **METHODS:** Bacterial binding and binding of monoclonal antibodies to H type 2, Lewis(a), and Lewis(b) to Kato III, buccal epithelial, and gastric mucosal cells were shown by flow cytometry. Bacterial outer membrane proteins eluted from H type 2, Lewis(a), or Lewis(b) were shown by polyacrylamide gel electrophoresis. **RESULTS:** Kato III and human epithelial cells bound each monoclonal antibody; O cells bound more anti-H type 2 ($P < 0.05$). Binding indices for *H. pylori* correlated with those for anti-H type 2 ($P < 0.005$) and anti-Lewis(b) ($P < 0.001$) but not anti-Lewis(a). A 61-kilodalton protein was eluted from H type 2, Lewis(a), or Lewis(b). **CONCLUSIONS:** Our results indicate that H type 2 is an important receptor for the 61-kilodalton bacterial adhesin, partly explaining increased susceptibility of individuals of blood group O to ulcers. Lewis(b) binds *H. pylori* more efficiently than Lewis(a). If these interactions occur in vivo, lack of Lewis(b) in mucosal fluids of nonsecretors may contribute to colonization by *H. pylori*.

PMID: 9098001 [PubMed - indexed for MEDLINE]

Request

□ 1: Gastroenterology. 1997 Apr;112(4):1179-87.

[Related Articles, Links](#)

Gastroenterology

Isolation of a cell surface component of *Helicobacter pylori* that binds H type 2, Lewis(a), and Lewis(b) antigens.

Alkout AM, Blackwell CC, Weir DM, Poxton IR, Elton RA, Luman W, Palmer K.

Department of Medical Microbiology, University of Edinburgh, Scotland.

BACKGROUND & AIMS: Individuals of blood group O and nonsecretors of ABO blood group antigens are more susceptible to peptic ulcers. The aim of this study was to determine if blood group antigens associated with group O or secretor status are epithelial cell receptors for *Helicobacter pylori*. **METHODS:** Bacterial binding and binding of monoclonal antibodies to H type 2, Lewis(a), and Lewis(b) to Kato III, buccal epithelial, and gastric mucosal cells were shown by flow cytometry. Bacterial outer membrane proteins eluted from H type 2, Lewis(a), or Lewis(b) were shown by polyacrylamide gel electrophoresis. **RESULTS:** Kato III and human epithelial cells bound each monoclonal antibody; O cells bound more anti-H type 2 ($P < 0.05$). Binding indices for *H. pylori* correlated with those for anti-H type 2 ($P < 0.005$) and anti-Lewis(b) ($P < 0.001$) but not anti-Lewis(a). A 61-kilodalton protein was eluted from H type 2, Lewis(a), or Lewis(b). **CONCLUSIONS:** Our results indicate that H type 2 is an important receptor for the 61-kilodalton bacterial adhesin, partly explaining increased susceptibility of individuals of blood group O to ulcers. Lewis(b) binds *H. pylori* more efficiently than Lewis(a). If these interactions occur in vivo, lack of Lewis(b) in mucosal fluids of nonsecretors may contribute to colonization by *H. pylori*.

PMID: 9098001 [PubMed - indexed for MEDLINE]



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[\[Sign In\]](#) [\[Register\]](#)

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search for

Limits Preview/Index History Clipboard Details

Display Show Sort by Send to

All: 1 Review: 0

About Entrez
NCBI Toolbar

Text Version

Entrez PubMed
Overview
Help | FAQ
Tutorials
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
Special Queries
LinkOut
My NCBI

Related Resources
Order Documents
NLM Mobile
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

☐ 1: [FEMS Immunol Med Microbiol.](#) 1996 Dec 1;16(2):141-55.

[Related Articles,](#)
[Links](#)

ELSEVIER
FULL-TEXT ARTICLE

Cell envelope characteristics of *Helicobacter pylori*: their role in adherence to mucosal surfaces and virulence.

Clyne M, Drumm B.

Department of Paediatrics, University College Dublin, Our Ladys Hospital for Sick Children, Crumlin, Ireland.

Helicobacter pylori colonises the gastric mucosa of humans and causes both antral gastritis and duodenal ulcer disease. Exactly how *H. pylori* causes disease is not known but several pathogenic determinants have been proposed for the organism. These include adhesins, cytotoxins and a range of different enzymes including urease, catalase and superoxide dismutase. Surface molecules of *H. pylori* such as flagella, lipopolysaccharide, the urease enzyme and outer membrane proteins are putative adhesin molecules. While phosphatidylethanolamine and the Lewis(b) blood group antigen have been proposed as receptor molecules for the organism the exact mechanism by which *H. pylori* adheres to the gastric mucosa has still to be identified. Characterisation of the adhesins of *H. pylori* could lead to the development of adhesin analogues for use in the inhibition of colonisation and improved therapy for ulcer disease. In vivo studies with isogenic mutants which are incapable of adhering to the gastric mucosa would greatly clarify the significance of adherence. Such mutants could possibly be useful as a vaccine against infection with wild-type organisms.

PMID: 8988394 [PubMed - indexed for MEDLINE]

Display Show Sort by Send to

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
Department of Health & Human Services
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Feb 13 2006 12:53:38

□ 1: APMIS. 1992 Oct;100(10):908-13.

[Related Articles](#), [Links](#)

Haemagglutination patterns of *Helicobacter pylori*. Frequency of sialic acid-specific and non-sialic acid-specific haemagglutinins.

Lelwala-Guruge J, Ljungh A, Wadstrom T.

Department of Medical Microbiology, University of Lund, Sweden.

Thirty-two *Helicobacter pylori* strains were screened for haemagglutination (HA) activity with erythrocytes of 11 different animal species. Twenty-three strains (72%) that agglutinated human erythrocytes exhibited a broad-spectrum HA profile. Human, guinea pig and bovine erythrocytes high in sialoglycoconjugates were strongly agglutinated by most strains. Except for two, seven strains (22%) that did not agglutinate human erythrocytes exhibited a narrow-spectrum HA profile, commonly not inhibitable by sialoglycoconjugates or N-acetylneuraminlactose (NANLac). Strains were classified into three major HA classes. HA of 10 strains (31%) in class I was inhibited by different combinations of NANLac, orosomucoid or fetuin, but not by asialofetuin, suggesting the presence of sialic acid-specific HAs probably recognizing NeuAc alpha-(2-3)-Gal isomer. Twelve strains (38%) in class II exhibited a different receptor specificity binding to different combinations of NANLac, orosomucoid and fetuin, as well as asialofetuin. No inhibition was observed with 10 strains (31%) in class III; thus, this receptor seems different from both the other classes. Of 21 strains (66%) in classes I and II, HA of 11 strains (34%) was inhibited with NANLac, 14 strains (44%) with orosomucoid and 15 strains (47%) with fetuin. The great heterogeneity observed in HA patterns indicates that the HAs of different strains may recognize a heterogeneous class of sialoglycoconjugates on the erythrocyte membrane.

PMID: 1445697 [PubMed - indexed for MEDLINE]

□ 1: J Infect. 1992 May;24(3):263-7.

[Related Articles, Links](#)

Adherence of *Helicobacter pylori* to HEp-2 cells.

Figueroa G, Portell DP, Soto V, Troncoso M.

Microbiology Unit, University of Chile, Santiago.

The adherence of 25 strains of *Helicobacter pylori* was evaluated in HEp-2 cells. These bacterial isolates, obtained from Chilean patients with gastric disorders, were also tested for haemagglutination of human red blood cells. Adherence of HEp-2 cells was expressed as a common property of all strains, irrespective of whether the cultures were grown on semi-solid or in liquid media. Previous reports that haemagglutinating activity was present in cells grown only on semi-solid media were confirmed. Adherence to HEp-2 cells was suppressed when bacterial cells were pretreated with homologous or heterologous whole human serum, containing specific antibodies of *H. pylori*. Adherence remained unaltered when bacterial cells were similarly treated with normal serum lacking specific antibodies. These observations imply that adhesions are expressed *in vivo* and suggest that an adherence mechanism, not depending on the expression of specific haemagglutinin antigen, operates for *H. pylori*.

PMID: 1602147 [PubMed - indexed for MEDLINE]

□ 1: FEMS Immunol Med Microbiol. 1995 Feb;10(3-4):307-16.

[Related Articles, Links](#)

Attachment, ingestion and intracellular killing of *Helicobacter pylori* by human peripheral blood mononuclear leukocytes and mouse peritoneal inflammatory macrophages.

Chmiela M, Paziak-Domanska B, Wadstrom T.

Department of Medical Microbiology, University of Lund, Sweden.

The different steps of phagocytosis, attachment, ingestion and intracellular killing of cells of *Helicobacter pylori* strain 17874 (expressing sialic acid-specific haemagglutinin) and cells of *H. pylori* strain 17875 (expressing non-sialic acid-specific haemagglutinin) have been studied. More cells of sialopositive *H. pylori* strain 17874 have been found attached to human peripheral blood mononuclear leukocytes (PBM) and mouse peritoneal inflammatory macrophages (PIM) than cells of sialonegative *H. pylori* strain 17875. Binding of cells of *H. pylori* strain 17874 has been significantly inhibited by treatment of phagocytes with neuraminidase. Inhibition of adhesion of these bacteria preincubated with foetuin to normal phagocytic cells has also been found. Well adhering cells of *H. pylori* strain 17874 were more resistant to killing mechanisms of human PBM and mouse PIM than cells of strain 17875. Good, probably sialic acid-specific haemagglutinin dependent, adhesion of *H. pylori* bacteria to phagocytes can be considered as an important virulence factor which facilitates the pathogen to avoid the defence mechanisms.

PMID: 7773248 [PubMed - indexed for MEDLINE]

[First Hit](#) [Fwd Refs](#)[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)

Generate Collection

Print

L1: Entry 11 of 15

File: USPT

Jun 25, 2002

US-PAT-NO: 6410719

DOCUMENT-IDENTIFIER: US 6410719 B1

TITLE: Blood group antigen binding protein and corresponding agents

DATE-ISSUED: June 25, 2002

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Boren; Thomas	S-906 28 Ume.ang.			SE
Arnqvist; Anna	Ume.ang.			SE
Normark; Staffan	Stockholm			SE
Ilver; Dag	Ume.ang.			SE

US-CL-CURRENT: [536/23.7](#); [536/23.1](#)

CLAIMS:

What is claimed is:

1. An isolated recombinant DNA from *Helicobacter pylori* encoding an adhesin protein, wherein said protein specifically binds to the fucosylated blood group antigens Lewis .sup.b and H-1, comprising the nucleotide acid sequence of SEQ ID NO:1.

2. A vector comprising an isolated recombinant DNA from *Helicobacter pylori* encoding an adhesin protein, wherein said protein specifically binds to the fucosylated blood group antigens Lewis.sup.b and H-1, comprising the nucleotide sequence of SEQ ID NO:1.

3. An isolated recombinant DNA from *Helicobacter pylori* encoding an adhesion protein, wherein said protein specifically binds to the fucosylated blood group antigens Lewis.sup.b and H-1, comprising the open reading frame of the nucleotide sequence of SEQ ID NO:1.

4. A vector comprising an isolated recombinant DNA from *Helicobacter pyroli* encoding an adhesin protein, wherein said protein specifically binds to the fucosylated blood group antigens Lewis.sup.b and H-1, comprising the open reading frame of the nucleotide sequence of SEQ ID NO:1.

[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)

[First Hit](#) [Fwd Refs](#) [Previous Doc](#) [Next Doc](#) [Go to Doc#](#)

[Generate Collection](#)[Print](#)

L1: Entry 9 of 15

File: USPT

Mar 23, 2004

US-PAT-NO: 6709656

DOCUMENT-IDENTIFIER: US 6709656 B1

TITLE: Helicobacter pylori adhesin binding group antigen

DATE-ISSUED: March 23, 2004

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Boren; Thomas	Umea			SE
Arnqvist; Anna	21 Umea			SE
Hammarstrom; Lennart	86 Huddinge			SE
Normark; Staffan	41 Stockholm			SE
Ilver; Dag	35 Umea			SE

US-CL-CURRENT: [424/190.1](#); [424/184.1](#), [424/234.1](#), [435/252.1](#), [435/7.32](#), [514/25](#),
[530/350](#)

CLAIMS:

What is claimed is:

1. An isolated and purified bacterial blood group antigen binding adhesin protein (BabA) from Helicobacter pylori species, wherein said protein binds specifically to fucosylated Lewis.sup.b type I and H-1 blood group antigen-glycoconjugates and, wherein said protein contains less than 20% bacterial protein impurities, has a molecular weight in the interval of 73 to 75 kDa as determined by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), and is not a HopA, HopB, HopC, HopD, or HopE protein.

2. The adhesion protein of claim 1, wherein the SEQ ID NO:5

EDDGFYTSVGYQIGEEAQMV

is in an amino terminal position.

3. The adhesion protein of claim 1, wherein the protein has a molecular weight of about 73.5 kDa, as determined by SDS-PAGE.

4. The adhesion protein of claim 3, wherein said Helicobacter pylori species is Helicobacter pylori strain CCUG 17875.

5. An immunogenic composition comprising an adhesion protein according to any one of claim 1, 2, 3, or 4, optionally together with pharmaceutically acceptable excipients.